arXiv:1702.05635v1 [math.CA] 18 Feb 2017

ON SOME INTEGRALS OF HARDY

ALEXANDER E PATKOWSKI

Abstract. We consider some properties of integrals considered by Hardy and Koshliakov, and which have also been further extended recently by Dixit. We establish a new general integral formula from some observations about the digamma function. We also obtain lower and upper bounds for Hardy's integral through properties of the digamma function.

Keywords: Fourier Integrals; Riemann xi function; Digamma function.
2010 Mathematics Subject Classification 11M06, 33C15.

1. Introduction and Main formulas

In a paper written by the well-known G. H. Hardy [10], an interesting integral formula is presented (corrected in [3])

(1.1)
$$\int_0^\infty \frac{\Xi(t/2)}{1+t^2}\frac{\cos(xt)}{\cosh(\pi t/2)}dt = \frac{1}{4}e^{-x}\left(2x+\frac{1}{2}\gamma+\frac{1}{2}\log(\pi)+\log(2)\right)+\frac{1}{2}e^x\int_0^\infty \psi(t+1)e^{-\pi t^2 e^{4x}}dt,$$

where $\psi(x) := \frac{\partial}{\partial x}\log(\Gamma(x))$, $\Gamma(x)$ being the gamma function [1], and $\Xi(t) := \xi(\frac{1}{2}+it)$, where [11, 15]

$$\xi(s) := \frac{1}{2}s(s-1)\pi^{-\frac{s}{2}}\Gamma(\frac{s}{2})\zeta(s).$$

Here we have used the standard notation for the Riemann zeta function $\zeta(s) := \sum_{n\geq 1} n^{-s}$, for $\Re(s) > 1$. Koshliakov [12] produced this formula as well, but in a slightly different form, using instead the integral

$$e^x\int_0^\infty (\psi(t+1)-\log(t))e^{-\pi t^2 e^{4x}}dt, \tag{1.2}$$

for the right side of (1.1). Here the trick, which has been exploited in the many studies [3, 4, 5, 6, 7, 12, 14], is to re-write the left side of (1.1) as an inverse Mellin transform by utilizing the classical functional equation $\xi(s) = \xi(1-s)$ [15]. Titchmarsh has used simpler but similar integrals as the left side of (1.1) to obtain Hardy's result that $\Xi(t)$ has infinitely many real zeros [15]. See Csordas [2] for some more recent work in this direction. A recent paper by Dixit offers a beautiful

generalization of formula (1.1), which involves a confluent hypergeometric function [3]. Other refinements can be found in [4, 5].

The purpose of this paper is to offer some further results concerning (1.1) that appear to have been overlooked. A new generalization is offered as well that takes a different route than the ones offered by [3, 4, 5], considering what is essentially the polygamma function [1]. In the following section, we offer some new inequalities that we noticed upon examining the integral given in (1.1).

Theorem 1.1. *For $m \in \mathbb{N}_0$, we have that*

$$(-1)^m e^x \int_0^\infty t^m \bar{\psi}_m(t) e^{-\pi t^2 e^{4x}} dt = \frac{1}{2}\int_0^\infty \frac{\Xi(t)}{\cosh(\pi t)(t^2+\frac{1}{4})}\left(\frac{\Gamma(\frac{1}{2}-it+m)}{\Gamma(\frac{1}{2}-it)}e^{-2xit} + \frac{\Gamma(\frac{1}{2}+it+m)}{\Gamma(\frac{1}{2}+it)}e^{2ixt}\right)dt,$$

$$\bar{\psi}_m(t) := \frac{\partial^m}{\partial t^m}(\psi(t+1)-\log(t)).$$

Proof. Recall the Mellin transform of a suitable function $f(t)$ is given by [1]

$$\int_0^\infty f(t)t^{s-1}dt = F(s),$$

and its inverse,

$$f(t) = \frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty} F(s)t^{-s}ds,$$

provided c is a real number chosen where $F(s)$ is analytic. This is a special application of Parseval's theorem for Mellin transforms with one function chosen as $f(t) = e^{-\pi t^2}$, and the other as $g(t) = t^m\bar{\psi}_m(t)$. From Titchmarsh [15, eq.(2.15.7)] we have for $0 < c < 1$,

$$\psi(x+1)-\log(x) = -\frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty}\frac{\pi\zeta(1-s)}{\sin(\pi s)}x^{-s}ds. \tag{1.3}$$

Differentiating (1.3) m times, gives us

$$\bar{\psi}_m(x) = (-1)^{m-1}\frac{1}{2\pi i}\int_{c-i\infty}^{c+i\infty}\frac{\pi\zeta(1-s)}{\sin(\pi s)}\frac{\Gamma(s+m)}{\Gamma(s)}x^{-s-m}ds. \tag{1.4}$$

Now proceeding as we mentioned in the beginning in the proof, noting $\sin((\frac{1}{2}+it)\pi) = \cosh(\pi t)$, the functional equation $\xi(s) = \xi(1-s)$, the Mellin transform of $e^{-\pi(xt)^2}$ is $\pi^{-s/2}x^{-s}\Gamma(s/2)/2$, with $0 < c' < 1$

$$\int_0^\infty f(t)g(t)dt = \frac{1}{2\pi i}\int_{c'-i\infty}^{c'+i\infty} F(s)G(1-s)ds,$$

we get $0 < c' < 1$

$$\frac{1}{2\pi i}\int_{c'-i\infty}^{c'+i\infty} \frac{\pi\xi(s)}{\sin(\pi s)}\frac{\Gamma(1-s+m)}{\Gamma(1-s)}x^{-s}ds.$$

Put $c' = \frac{1}{2}$ and we find this is equal to

$$\frac{1}{2\sqrt{x}}\int_{-\infty}^{\infty} \frac{\Xi(t)}{\cosh \pi t}\frac{\Gamma(\frac{1}{2}-it+m)}{\Gamma(\frac{1}{2}-it)}x^{-it}dt.$$

Splitting this bilateral integral up into two integrals and replacing x by e^{2x} leads to the result.

□

Koshliakov [12] gives a similar integral to (1.1) but squaring the $\Xi(t)(t^2+\frac{1}{4})^{-1}$ function. Moll and Dixit state his result in [6, Theorem 4.4], and offer further interesting generalizations. However, we noticed a slightly different form of this integral based on the type of proof we used in the previous theorem.

Theorem 1.2. *For $x > 0$,*

(1.5)

$$\int_0^\infty (\psi(t+1)-\log(t))\left(\frac{2\pi e^{x/2}}{t} - e^{-x/2}4\sum_{n\geq 1}K_0(2\pi n t e^{-x})\right)dt = \int_0^\infty \frac{\Xi^2(t)}{(t^2+\frac{1}{4})^2}\frac{\cos(xt)}{\cosh(\pi t)}dt.$$

Proof. First, since $K_0(t) = O(t^{-1/2}e^{-t})$, for $t \to \infty$ in $|\arg(t)| < 3\pi/2$ where as usual $K_s(t)$ is the modified Bessel function, it follows that the series on the right side of the formula [9] $d > 0$,

$$\sum_{n\geq 1}\frac{1}{2\pi i}\int_{d-i\infty}^{d+i\infty}\Gamma^2(\frac{s}{2})(\frac{x}{2n})^{-s}ds = 4\sum_{n\geq 1}K_0(n\pi x),$$

converges absolutely. Summing through, we find for $d' > 1$,

$$\frac{1}{2\pi i}\int_{d'-i\infty}^{d'+i\infty}\Gamma^2(\frac{s}{2})(\frac{x}{2})^{-s}\zeta(s)ds = 4\sum_{n\geq 1}K_0(n\pi x),$$

which has a simple pole at $s = 1$. If we calculate this residue and move the line of integration to $\Re(s) = b$, $0 < b < 1$, we find

$$\frac{1}{2\pi i}\int_{b-i\infty}^{b+i\infty}\Gamma^2(\frac{s}{2})(\frac{x}{2})^{-s}\zeta(s)ds + \frac{2\pi}{x} = 4\sum_{n\geq 1}K_0(n\pi x). \tag{1.6}$$

Now noting (1.3) and (1.6) in conjunction with Parseval's theorem for Mellin transforms, we obtain the result after applying the functional equation for $\xi(s)$. □

Other associated integral formulae related to Fourier cosine transforms may be obtained from using known evaluations.

Theorem 1.3. *We have for* $\frac{\pi}{2} > |\Im(\beta')|$,

$$\int_0^\infty \frac{\cosh(t)}{\cosh(2t)+\cosh(2\beta')}\left(e^{t/2}-2e^{-t/2}\sum_{n\geq 1}e^{-\pi n^2 e^{-2t}}\right)dt \tag{1.7}$$

$$=\frac{\pi e^{\beta'}}{4\cosh(\beta')}\int_0^\infty (\psi(x+1)-\log(x))e^{-\pi x^2 e^{4\beta'}}\,dx.$$

Proof. We apply the classical [15, eq.(2.16.2)]

$$\int_0^\infty \frac{\Xi(t)}{(t^2+\frac{1}{4})}\cos(xt)dt=\frac{\pi}{2}(e^{t/2}-2e^{-t/2}\sum_{n\geq 1}e^{-\pi n^2 e^{-2t}}),$$

with the formula [9], valid for $\pi\Re(\alpha) > |\Im(\bar{\alpha}\beta)|$, and

$$\int_0^\infty \frac{\cos(ty)\cosh(t\alpha/2)dt}{\cosh(\alpha t)+\cosh(\beta)}=\frac{\pi}{2\alpha}\frac{\cos(\beta y/\alpha)}{\cosh(\beta/2)\cosh(\pi y/\alpha)}.$$

We need only let $\beta \to 2\beta'$, and $\alpha = 2$, and then we have $\frac{\pi}{2} > |\Im(\beta')|$. This coupled with Parseval's theorem gives the result.

□

Note that the integral on the left side of (1.6) hints to us about the non-negativity of the integral on the right side. We will investigate this further in the next section.

2. Inequalities

We first start with some preliminaries about some special functions which can be found in [1]. First we recall the error function,

$$\operatorname{erf}(x)=\frac{2}{\sqrt{\pi}}\int_0^x e^{-t^2}dt,$$

and its complement, $\operatorname{erfc}(x) = 1-\operatorname{erf}(x)$. Also, $\operatorname{erfi}(x) := -i\operatorname{erf}(ix)$. We also use the exponential integral

$$\operatorname{Ei}(x)=-\int_{-x}^\infty \frac{e^{-t}}{t}dt.$$

Theorem 2.1. *Define the Hardy integral for* $y > 0$ *as* $I(y) := \int_0^\infty(\psi(t+1)-\log(t))e^{-yt^2}dt$. *Then we have,*

$$\frac{(2c_1)^2\sqrt{y}}{\sqrt{\pi}\operatorname{erfi}(\sqrt{y})}-\frac{\operatorname{Ei}(-y)}{4}-\frac{e^{-y}}{12}+\frac{\sqrt{y\pi}}{12}\operatorname{erfi}(\sqrt{y})\leq I(y).$$

Furthermore,

$$I(y) \leq \sqrt{c_2\sqrt{\frac{\pi}{2y}}\operatorname{erf}(\sqrt{2y})} - \frac{\operatorname{Ei}(-y)}{4},$$

where $c_1 = 0.952894$, *and* $c_2 = 1.56624$.

Proof. We first need a result from the paper [13, Corollary 1], for $x > 0$,

$$\frac{1}{2x} - \frac{1}{12x^2} < \psi(x+1) - \log(x) < \frac{1}{2x}. \tag{2.1}$$

For the upper bound we first write,

$$I(y) = \int_0^1 (\psi(t+1) - \log(t))e^{-yt^2}dt + \int_1^\infty (\psi(t+1) - \log(t))e^{-yt^2}dt.$$

Applying (2.1) we have that this is

$$\leq \int_0^1 (\psi(t+1) - \log(t))e^{-yt^2}dt - \frac{\operatorname{Ei}(-y)}{4}.$$

By Schwarz's inequality and the fact that

$$\int_0^1 (\psi(t+1) - \log(t))^2 dt = 1.56624$$

we have the upper bound. For the lower bound, we write the Schwarz's inequality in the form

$$\left(\int_0^1 (\psi(t+1) - \log(t))^{1/2}dt\right)^2 \leq \left(\int_0^1 (\psi(t+1) - \log(t))e^{-yt^2}dt\right)\left(\int_0^1 e^{yt^2}dt\right),$$

and also note,

$$\int_0^1 \sqrt{(\psi(t+1) - \log(t))}dt = 0.952894.$$

This coupled with similar computations derived from the left side of (2.1), together with the definitions of the special integral functions, gives our result. □

Now it can be seem that the lower bound actually implies the Hardy integral $I(x)$ is non-negative for $x > 0$, and also the squeeze theorem shows that $I(x) \to 0$ when $x \to \infty$.

3. Other observations and further comments

While studies such as this one and others have shown there is a plethora of integral formulae like Hardy's (1.1) to be found, it would be interesting to see if these integrals could be incorporated into similar work like Csordas [2]. We were interested in seeing if (1.1) could be adapted to imply information beyond Hardy's theorem, that $\Xi(t)$ has infinitely many real zeros. The classical ideas are included in Titchmarsh's text [15, pg. 256-260], and evolve around the ideas of first assuming $\Xi(t)$ is of one sign, and then arriving at a contradiction from properties of an integral of the form $\int_0^\infty \Xi(t)k(t)dt$, $k(t)$ non-negative for $t > 0$. The idea in [15, pg. 260] considers the moments $k(t) = t^{2n}$. Apply the operator $\bar{\partial}_x = -\partial_x^2 + \frac{1}{4}$ of Glasser [8] to (1.1) (after replacing t by $t/2$) and set $x = x' + i\pi$ to get

$$\lim_{x \to x'+i\pi} \bar{\partial}_x(e^x I(\pi e^{4x})) = \int_0^\infty \Xi(t)\cos(x't)dt + i\int_0^\infty \Xi(t)\frac{\sin(x't)\sinh(\pi t)}{\cosh(\pi t)}dt.$$

Differentiating $2n$ times and setting $x' = 0$ gives us the right side as the value $(-1)^n \int_0^\infty t^{2n}\Xi(t)dt$. Note that $e^{x'+i\pi}I(\pi e^{4(x'+i\pi)}) = -e^{x'}I(\pi e^{4x'})$. Further the integral $I(x)$ clearly converges only for $\Re(x) > 0$, and hence $I(\pi e^{4z})$ converges when $-\frac{\pi}{4} < \Im(z) < \frac{\pi}{4}$, and $\lim_{z \to i\pi/4} I(\pi e^{4z}) = +\infty$. It would be interesting if new information could be obtained about $\Xi(t)$ from alternative forms of these moments like those found here.

References

[1] G. Andrews, R. Askey, and R. Roy. *Special Functions,* volume 71 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, New York, 1999.

[2] G. Csordas, *Fourier Transforms of Positive Definite Kernels and the Riemann ξ-function,* Computational Methods and Function Theory, Volume 15, Issue 3, pp 373–391 (2015).

[3] A. Dixit. *Analogues of the general theta transformation formula,* Proc. Roy. Soc. Edinburgh, Sect. A, 143:371–399, 2013.

[4] A. Dixit, N. Robles, A. Roy and A. Zaharescu, *Zeros of combinations of the Riemann ξ-function on bounded vertical shifts,* J. Number Theory Volume 149, April 2015, Pages 404?434

[5] A. Dixit, A. Roy and A. Zaharescu, *Riesz-type criteria and theta transformation analogues,* J. Number Theory 160, p. 385–408 (2016).

[6] A. Dixit, V. H. Moll *Self-reciprocal functions, powers of the Riemann zeta function and modular-type transformations,* J. Number Theory 147 (2015), 211–249.

[7] W. L. Ferrar, *Some solutions of the equation $F(t) = F(t^{-1})$,* J. London Math. Soc. 11 (1936), 99–103.

[8] M.L. Glasser *A note on the Fourier transform,* Journal of Physics A: Mathematical and Theoretical, Volume 49, Number 45 (2016), 454004.

[9] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products.* Edited by A.Jeffrey and D. Zwillinger. Academic Press, New York, 7th edition, 2007.

[10] G.H. Hardy, *Note by Mr. G.H. Hardy on the preceding paper,* Quart. J. Math. 46 (1915), 260–261.

[11] H. Iwaniec and E. Kowalski, *Analytic number theory,* American Mathematical Society Colloquium Publications, vol. 53, American Mathematical Society, Providence, RI, 2004.

[12] N. S. Koshlyakov, *On a transformation of definite integrals and its application to the theory of Riemann's function* $\zeta(s)$*,* Comp. Rend. (Doklady) Acad. Sci. URSS 15 (1937), 3–8.

[13] Feng Qi, Run-Qing Cui, Chao-Ping Chen, Bai-Ni Guo *Some completely monotonic functions involving polygamma functions and an application,* J. Math. Anal. Appl. 310 (2005) 303–308.

[14] S. Ramanujan, *New expressions for Riemann's functions* $\xi(s)$ *and* $\Xi(t)$*,* Quart. J. Math., 46: 253–260, 1915.

[15] E. C. Titchmarsh, *The theory of the Riemann zeta function,* Oxford University Press, 2nd edition, 1986.

1390 Bumps River Rd.

Centerville, MA 02632

USA

E-mail: alexpatk@hotmail.com